



Utah

Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
38th highest	33rd highest	35th highest

*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




Sexually transmitted diseases Utah

Gonorrhea rates, 1982–1992



Gonorrhea rates, 1992



National ranking,* 1992


Males – 43rd highest Females – 45th highest

**National ranking refers to males and females aged 10–24 years.*




Chlamydia cases and rates,* 1988–1992


Cases




Rates




Cases



Rates



Acquired immunodeficiency syndrome (AIDS) cumulative cases,† 1981–1992



*Chlamydia cases and rates refer to males and females of all ages.


†Missing bar indicates fewer than six cumulative cases.




Related risk behaviors,* 1993

Utah


Had sexual intercourse (lifetime)




Had sexual intercourse during the past three months (currently sexually active)




Used birth control pills during last sexual intercourse (among currently sexually active)



Had sexual intercourse with four or more people (lifetime)



Used a condom during last sexual intercourse (among currently sexually active)



Injected illegal drugs (lifetime)



*Related risk behaviors refer to high school students.

Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 Utah Youth Risk Behavior Survey (risk behavior data).

